

**Influenza A H1N1 (A/California/04/2009)  
Hemagglutinin / HA Antibody, Rabbit PAb,  
Antigen Affinity Purified**

Catalog Number: 11055-T44



**Sino Biological**  
Biological Solution Specialist

GENERAL INFORMATION	
<b>Immunogen:</b>	Recombinant Influenza A H1N1 (A/California/04/2009) Hemagglutinin / HA protein (Catalog#11055-V08H)
<b>Preparation</b>	Produced in rabbits immunized with purified, recombinant Influenza A H1N1 (A/California/04/2009) Hemagglutinin / HA (Catalog#11055-V08H; ACP41105.1; Met1-Gln529). Influenza A H1N1 (A/California/04/2009) Hemagglutinin / HA specific IgG was purified by Influenza A H1N1 (A/California/04/2009) Hemagglutinin / HA affinity chromatography.
<b>Ig Type:</b>	Rabbit IgG
<b>Specificity:</b>	Influenza A H1N1 (A/California/04/2009) Hemagglutinin / HA
<b>Formulation:</b>	0.2 µm filtered solution in PBS
<b>Storage:</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.
APPLICATIONS	
<b>Applications:</b>	WB, IHC-P, IP
	(Antibody's applications have not been validated with corresponding viruses. Optimal concentrations/dilutions should be determined by the end user.)
RECOMMENDED CONCENTRATION	
<b>Western Blot</b>	WB: 1:500-1:2000


**Please Note: Optimal concentrations/dilutions should be determined by the end user.**

**Influenza A H1N1 (A/California/04/2009)  
Hemagglutinin / HA Antibody, Rabbit PAb,  
Antigen Affinity Purified**

Catalog Number: 11055-T44



Sino Biological  
Biological Solution Specialist



Anti-Influenza A H1N1 (A/California/04/2009)  
Hemagglutinin / HA rabbit polyclonal antibody  
at 1:500 dilution.

Sample: Influenza A H1N1  
(A/California/04/2009) Hemagglutinin / HA  
Recombinant Protein

Lane A: 100ng

Lane B: 10ng

Secondary

Goat anti Rabbit IgG (H+L)/HRP at 0.4ug/ml

Performed under reducing conditions.