

Influenza A H5N8 (A/duck/Jiangsu/k1203/2010) Hemagglutinin / HA1 HEK293 Cell Lysate (WB positive control)

Catalog Number: 40385-V08H1L



Sino Biological Inc.
Biological Solution Specialist

Hemagglutinin / HA Transfected / Overexpression Cell Lysate Product Information

Expressed Host:	HEK293 Cells
Products Description:	Human Cell Lysate that Influenza A H5N8 (A/duck/Jiangsu/k1203/2010) Hemagglutinin / HA1 transfected / overexpressed for Western blot (WB) positive control. The whole cell lysate is provided in 1X Sample Buffer (1X modified RIPA buffer+1X SDS loading buffer).
Sequence information:	A DNA sequence encoding the Influenza A virus (A/duck/Jiangsu/k1203/2010(H5N8)) hemagglutinin (AFK13933.1) (Met1-Arg345), termed as HA1, was expressed with a polyhistidine tag at the C-terminus.
Predicted N Terminal:	Asp 17
Molecule Mass:	The recombinant HA1 subunit of the Influenza A virus (A/duck/Jiangsu/k1203/2010(H5N8)) hemagglutinin consists of 340 amino acids and predicts a molecular mass of 38.7 kDa.
Species:	H5N8

Hemagglutinin / HA Transfected / Overexpression Cell Lysate Usage Guide

Preparation Method:	Cell lysate was prepared by homogenization in ice-cold modified RIPA Lysis Buffer with cocktail of protease inhibitors (Sigma). Cell debris was removed by centrifugation. Protein concentration was determined by Bradford assay (Bio-Rad protein assay, Microplate Standard assay). The cell lysate was boiled for 5 min in 1 x SDS loading buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis Buffer:	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF.
Quality Control Testing:	12.5% SDS-PAGE Stained with Coomassie Blue after protein purification.
Stability:	Samples are stable for up to twelve months from date of receipt.
Recommend Usage:	<ol style="list-style-type: none">1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200µL pure water and boil for 2-5 min.3. Store the lyophilized cell lysate at 4°C. After re-dissolution, recommend to aliquot it into smaller quantities and store at -80°C.
Storage Buffer:	1 X Sample Buffer (1 X modified RIPA buffer+1 X SDS loading buffer).
Storage Instruction:	Store at 4°C. After re-dissolution, aliquot and store at -80°C.
Application notes:	Western blot (WB): Use at an assay dependent dilution. Other Applications: Not tested. Optimal dilutions/concentrations should be determined by the end user.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217

● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288

● Tel:+86-400-890-9989 ●

● <http://www.sinobiological.com>