

# Influenza A H1N1 (A/Texas/36/1991) Hemagglutinin / HA Insect Cell Lysate (WB positive control)

Catalog Number: 40132-V08BL



**Sino Biological Inc.**  
Biological Solution Specialist

## Hemagglutinin / HA Transfected / Overexpression Cell Lysate Product Information

|                              |  |
|------------------------------|--|
| <b>Expressed Host:</b>       | Baculovirus-Insect Cells   |
| <b>Products Description:</b> | Baculovirus-Insect Cell lysate that Influenza A H1N1 (A/Texas/36/1991) Hemagglutinin / HA transfected / overexpressed for Western blot (WB) positive control. The whole cell lysate is provided in 1X Sample Buffer (1X modified RIPA buffer+1X SDS loading buffer). |
| <b>Sequence information:</b> | A DNA sequence encoding the Influenza A virus (A/Texas/36/1991(H1N1)) hemagglutinin (ACF41933.1) (Met1-Gln529), termed as HA, was expressed with a polyhistidine tag at the C-terminus.  |
| <b>Predicted N Terminal:</b> | Asp 18   |
| <b>Molecule Mass:</b>        | The recombinant hemagglutinin of Influenza A virus (A/Texas/36/1991(H1N1)) consists 523 amino acids and predicts a molecular mass of 59.1 kDa.   |
| <b>Species:</b>              | H1N1   |

## Hemagglutinin / HA Transfected / Overexpression Cell Lysate Usage Guide

|                                 |  |
|---------------------------------|--|
| <b>Preparation Method:</b>      | Cell lysate was prepared by homogenization in ice-cold modified RIPA Lysis Buffer with cocktail of protease inhibitors (Sigma). Cell debris was removed by centrifugation. Protein concentration was determined by Bradford assay (Bio-Rad protein assay, Microplate Standard assay). The cell lysate was boiled for 5 min in 1 x SDS loading buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized. |
| <b>Lysis Buffer:</b>            | Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF.   |
| <b>Quality Control Testing:</b> | 12.5% SDS-PAGE Stained with Coomassie Blue after protein purification.   |
| <b>Stability:</b>               | Samples are stable for up to twelve months from date of receipt.   |
| <b>Recommend Usage:</b>         | <ol style="list-style-type: none"><li>1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.</li><li>2. Re-dissolve the pellet using 200µL pure water and boil for 2-5 min.</li><li>3. Store the lyophilized cell lysate at 4°C. After re-dissolution, recommend to aliquot it into smaller quantities and store at -80°C.</li></ol>  |
| <b>Storage Buffer:</b>          | 1 X Sample Buffer (1 X modified RIPA buffer+1 X SDS loading buffer).   |
| <b>Storage Instruction:</b>     | Store at 4°C. After re-dissolution, aliquot and store at -80°C.  |
| <b>Application notes:</b>       | Western blot (WB): Use at an assay dependent dilution.<br>Other Applications: Not tested.<br>Optimal dilutions/concentrations should be determined by the end user.  |

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217

● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288

● Tel:+86-400-890-9989 ●

● <http://www.sinobiological.com>